

ACTHAR

Products Affected

- HP ACTHAR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple sclerosis AND patient is currently on a disease modifying drug (interferon beta 1a, peginterferon beta1a, interferon beta1b, glatiramer, natalizumab, mitoxatrone, dimethyl fumarate, teriflunomide, alemtuzumab) to control disease progression, OR has tried and failed, contraindicated or intolerant to all DMDs AND the patient has failed corticosteroid therapy within the last 30 days or has an FDA labeled contraindication to corticosteroid therapy.
Age Restrictions	Infantile spasms patient must be less than 24 months of age
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form. Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline). Pregnancy.
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH .Chronic thromboembolic pulmonary hypertension (CTEPH) (Initial): One of the following: A) Both of the following: 1) Diagnosis of inoperable or persistent/recurrent CTEPH and 2) CTEPH is symptomatic OR B) Patient is currently on any therapy for the diagnosis of CTEPH. For all indications female patients are enrolled in the ADEMPAS REMS program.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	None

AFINITOR

Products Affected

- AFINITOR
- AFINITOR DISPERZ

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with Sutent or Nexavar OR Diagnosis of pancreatic neuroendocrine tumors (pNET) that are unresectable, locally advanced, or metastatic OR Diagnosis of renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery OR Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin OR Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection OR diagnosis of adult patients with progressive, well-differentiated, non-functional, neuroendocrine tumors (NET) of gastrointestinal (GI) or lung origin with unresectable, locally advanced, or metastatic disease OR diagnosis of tuberous sclerosis complex (TSC) associated partial seizures.
Age Restrictions	18 years of age and older for RCC, pNET, and renal angiomyolipoma with TSC. 1 year of age and older for SEGA. 2 years of age and older for TSC associated partial seizures.
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ALECENSA

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase(ALK) positive non-small cell lung cancer detected with an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND a history of failure, contraindication, intolerance, or progressed on XALKORI (crizotinib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ALUNBRIG

Products Affected

- ALUNBRIG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, ALK positive non-small cell lung cancer and have progressed or are intolerant to Xalkori (crizotinib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

AMPYRA

Products Affected

- AMPYRA
- *dalfampridine er*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of seizure. Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute).
Required Medical Information	Diagnosis of multiple sclerosis. Patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting Ampyra and patient is currently on a disease modifying drug (interferon beta 1a, peginterferon beta 1a, interferon beta 1b, or glatiramer) to control disease progression, or has documented treatment failure, intolerance, or contraindication to any one of the following: interferon beta 1a, peginterferon beta 1a, interferon beta 1b, or glatiramer.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Initial: 3 months, Reauthorization: 12 months
Other Criteria	None

AUBAGIO

Products Affected

- AUBAGIO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severe hepatic impairment. Current treatment with leflunomide. Patients who are pregnant or women of childbearing potential not using reliable contraception.
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS or progressive-relapsing MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis AND Patient has tried and had an insufficient response to at least one other formulary MS disease modifying therapy (e.g., Avonex, Betaseron, Copaxone, Gilenya, Tecfidera)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)

AURYXIA

Products Affected

- AURYXIA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	For the management of hyperphosphatemia in patients with chronic kidney disease on dialysis
Age Restrictions	18 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

BEXAROTENE

Products Affected

- *bexarotene*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient must meet one of following criteria: received prior systemic therapy for CTCL OR advanced-stage MF (stage IIB, III or IV) or SS OR early-stage MF (stage IA, IB or IIA) with folliculotropic/large cell transformation OR early-stage MF (stage IA, IB or IIA) refractory to skin directed therapy.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	Definite diagnosis of cutaneous T-cell lymphoma (CTCL) AND refractory to any prior systemic therapy (such as methotrexate)

BOSULIF

Products Affected

- BOSULIF

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Philadelphia chromosome-positive (Ph+) CML with resistance, relapse, or inadequate response to prior therapy with one of the following tyrosine kinase inhibitors (TKI): Gleevec [imatinib] OR Tasigna [nilotinib] OR newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (Ph + CML)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

BRIVIACT

Products Affected

- BRIVIACT ORAL

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of partial-onset seizures, member must have history of inadequate response, contraindication, or intolerance to levetiracetam prior to approval.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

CABOMETYX

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients who have or are at risk for severe hemorrhage and/or patients with a recent history of bleeding or hemoptysis.
Required Medical Information	Diagnosis of advanced renal cell carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

CALQUENCE

Products Affected

- CALQUENCE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	MANTLE CELL LYMPHOMA (MCL) (1) Patient must have a diagnosis of MCL AND (2) Patient has tried one other therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

CAPRELSA

Products Affected

- CAPRELSA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of medullary thyroid cancer (MTC), and disease is one of the following: A) unresectable, locally advanced, or B) metastatic AND one of the following: patient has symptomatic disease or patient has progressive disease.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

CARIMUNE

Products Affected

- CARIMUNE NF INTRAVENOUS SOLUTION RECONSTITUTED 6 GM

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Diagnosis of a primary humoral immunodeficiency disorder such as: primary immunoglobulin deficiency syndrome X-linked immunodeficiency with hyperimmunoglobulin etc). Documented hypogammaglobulinemia (IgG less than 600mg/dl) Idiopathic/Immune Thrombocytopenia Purpura. Diagnosis of Acute ITP with any of the following: Management of acute bleeding due to severe thrombocytopenia (platelets less than 30 000/mcL). To increase platelet counts prior major surgical procedures. Severe thrombocytopenia (platelets less than 20 000/mcL) at risk for intracerebral hemorrhage. Diagnosis of Chronic ITP and ALL of the following are met: Prior treatment has included corticosteroids and splenectomy Duration of illness less than 6 months, no concurrent illness explaining thrombocytopenia, platelets persistently at or below 20 000/mcL. Chronic Lymphocytic Leukemia (CLL B-cell) with either of the following present: Hypogammaglobulinemia (IgG less than 600mg/dL) or Recurrent bacterial infections associated with B-cell CLL. Kawasaki Disease-Diagnosed with Kawasaki Syndrome within ten days of onset of disease manifestations or is diagnosed after ten days of disease onset and continues to exhibit manifestations of inflammation or evolving coronary artery disease. IVIG is used in combination with high dose aspirin for the prevention of coronary artery aneurysms. Bone Marrow Transplant (BMT). Member is hypogammaglobinemic (IgG</p>

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
 Last Updated: 12/12/2018
 Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
	less than 400mg/dL). Hematopoietic Stem Cell Transplantation (HSCT). Is within first 100 days of allogenic hematopoietic stem cell transplantation. Is experiencing hypogammaglobulinemia (serum IgG level less than 400 mg/dL). AIDS/HIV- Has any of the following conditions: CD4+ T-cell counts greater than or equal 200/mm ³ , to prevent maternal transmission of HIV infection, IVIG is used in conjunction with zidovudine to prevent serious bacterial infections in HIV-infected members who have hypogammaglobulinemia (serum IgG less than 400 mg/dL).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

COMETRIQ

Products Affected

- COMETRIQ (100 MG DAILY DOSE)
- COMETRIQ (140 MG DAILY DOSE)
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Gastrointestinal perforation. Fistula. Severe hemorrhage.
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

COPAXONE

Products Affected

- COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- *glatiramer acetate*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsing-remitting multiple sclerosis OR diagnosis of first clinical episode with MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient does not have progressive disease and responding to therapy.

COTELLIC

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable OR metastatic malignant melanoma with BRAF V600E OR V600K mutation. Documentation of combination therapy with vemurafenib (Zelboraf)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

DICLOFENAC TOPICAL

Products Affected

- *diclofenac sodium transdermal gel 1 %, 3 %*
- *diclofenac sodium transdermal solution*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diclofenac 1% gel or 1.5% solution: Diagnosis of osteoarthritis, diclofenac 3% gel: Diagnosis of actinic keratosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ENDARI

Products Affected

- ENDARI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute sickle cell disease AND Must have A) trial history of Hydroxyurea OR B) intolerance to Hydroxyurea OR C) contraindication to Hydroxyurea
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ERLEADA

Products Affected

- ERLEADA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of non-metastatic, castration-resistant prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ESBRIET

Products Affected

- ESBRIET

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	1)Diagnosis of Idiopathic pulmonary fibrosis (IPF) as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity), AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF, AND 2) not used in combination with Ofev (nintedanib).
Age Restrictions	None
Prescriber Restrictions	Pulmonologist
Coverage Duration	12 months
Other Criteria	None

ESRD THERAPY

Products Affected

- PROCRIT

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Epoetin alfa therapy is not considered medically necessary for members with the following concomitant conditions: Concomitant use of another Recombinant Erythropoietin Product, anemia in cancer not related to chemotherapy OR anemia associated only with radiotherapy (without chemo). ESAs are not indicated in cancer patients receiving myelosuppressive chemotherapy when the anticipated outcome is cure (ESAs remain indicated when myelosuppressive chemotherapy is intended for palliation).
Required Medical Information	Pretreatment hemoglobin levels of less than 10g/dL. Dose reduction or interruption if hemoglobin exceeds 10 g/dL (CKD not on dialysis-adult, cancer), 11 g/dL (CKD on dialysis), 12 g/dL (pediatric CKD) in addition to supporting statement of diagnosis from physician.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None

FARYDAK

Products Affected

- FARYDAK

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Multiple Myeloma (MM) Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

FENTORA

Products Affected

- FENTORA BUCCAL TABLET 100 MCG
- LAZANDA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Fentanyl Buccal Tablet (Fentora) may be considered medically necessary when the following criteria are met: The member is currently diagnosed with cancer, OTFC is required to manage break-through pain, member is currently taking opioid therapy and is opioid tolerant, and member has tried and failed generic Actiq (fentanyl citrate lozenge). Tolerance is defined as any of the following: greater than or equal to 60mg morphine/day for greater than or equal than 1 week or greater than or equal to 50 mcg transdermal fentanyl/hour for greater than or equal 1 week or an equi-analgesic dose of another opioid for greater than or equal 1 week.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

FORTEO

Products Affected

- FORTEO SUBCUTANEOUS SOLUTION 600 MCG/2.4ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient has a diagnosis of one of the following: a) osteoporosis in a postmenopausal female, b) primary or hypogonadal osteoporosis in a male, or c) osteoporosis associated with sustained systemic glucocorticoid therapy AND patient is considered to be at high-risk for fracture by meeting one or more of the following: A) history of osteoporotic fracture, B) Low Bone Density less than 2.5 SD below normal, AND one or more of the following: i) failed one oral bisphosphonate and 1 injectable bisphosphonate, or ii) intolerant to one oral bisphosphonate and one injectable bisphosphonate. Patient has not received more than 2 years of therapy with Forteo.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Approve doses based on FDA labeling

GILENYA

Products Affected

- GILENYA ORAL CAPSULE 0.5 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure. History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker. Baseline QTC interval greater than or equal to 500 milliseconds. Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol).
Required Medical Information	Diagnosis of a relapsing form of multiple sclerosis or diagnosis of first clinical episode with MRI features consistent with MS AND Patient will be observed for signs and symptoms of bradycardia in a controlled setting for at least 6 hours after the first dose
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	For renewal, the patient has experienced no or slowed disease progression.

GILOTRIF

Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician in patients with: 1) metastatic non-small cell lung cancer (NSCLC) whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test or 2) metastatic squamous NSCLC, progressing after platinum-based chemotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

GOCOVRI

Products Affected

- GOCOVRI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients with ESRD (CrCl below 15 ml/min/m ²)
Required Medical Information	INITIAL: Diagnosis of Parkinsons disease AND (1) Patient is experiencing dyskinesia AND (2) Patient is receiving levodopa based therapy AND (3) Must have documented trial and failure to amantadine immediate release. RENEWAL: (1) must meet the initial criteria above AND (2) Documentation of positive clinical response to Gocovri (e.g., decreased off periods, decreased on time with troublesome dyskinesia)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None

GROWTH HORMONE

Products Affected

- NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Acute critical illness due to complications following open heart or abdominal surgery, multiple accidental trauma or acute respiratory failure. Active malignancy. Active proliferative or severe non-proliferative diabetic retinopathy. For PWS only: severe obesity, history of upper airway obstruction or sleep apnea, or severe respiratory impairment.
Required Medical Information	Diagnosis of pediatric indication: A) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B) SGA and birth weight or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SDS below mean for age and gender), C) CRI and nutritional status has been optimized, metabolic abnormalities have been corrected, and patient has not had renal transplant D) SHOX deficiency or Noonan syndrome E) PWS confirmed by genetic testing, F) Turner Syndrome confirmed by chromosome analysis. For GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following: height more than 3 SDS below mean for age and gender, or height more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of an adult indication: A) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
	<p>contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy) AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications AND Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH).</p>
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HEPATITIS B

Products Affected

- *adefovir dipivoxil*
- *entecavir*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients that have immune-tolerant chronic hepatitis B per AASLD guidelines
Required Medical Information	Must submit documentation of immune-active chronic hepatitis B per AASLD guidelines.
Age Restrictions	None
Prescriber Restrictions	Prescriber must be a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None

HEPATITIS C

Products Affected

- MAVYRET
- VOSEVI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Must submit documentation of chronic hepatitis C genotype (confirmed by HCV RNA level within the last 6 months). Must submit laboratory results within 6 weeks of initiating therapy including: 1) CBC w Platelets, 2) AST/ALT, 3) Total Bilirubin, 4) Serum Albumin, 5) PT/INR, 6) Serum Creatinine, and 7) GFR. FOR all GENOTYPES-Trial/failure, contraindication to, or intolerance to Mavyret required prior to the approval of Vosevi.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescriber must be a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None

HETLIOZ

Products Affected

- HETLIOZ

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Non-24-hour-sleep-wake disorder (Non-24) AND patient has documented blindness
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	For renewal, patient experienced an objective improvement (e.g., improvement in timing of nighttime sleep, improvement in duration of nighttime sleep, or reduction in daytime sleep).

HRM

Products Affected

- *guanfacine hcl er*
- *methylphenidate hcl er (cd) oral capsule extended release 10 mg, 20 mg, 40 mg, 50 mg, 60 mg*
- *methylphenidate hcl er oral tablet extended release 20 mg*
- *methylphenidate hcl er oral tablet extended release 24 hour*
- *methylphenidate hcl oral*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk. AND The physician must document on progress notes that patient counseling has and will continue to take place outlining the risks and potential side effects of the medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-ANALGESICS

Products Affected

- ASCOMP-CODEINE
- *butalbital-acetaminophen oral tablet 50-325 mg*
- *butalbital-apap-caff-cod*
- *butalbital-apap-caffeine oral capsule*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *indomethacin er*
- *indomethacin oral*
- *ketorolac tromethamine oral*
- *meperidine hcl injection solution 100 mg/ml, 25 mg/ml, 50 mg/ml*
- *meperidine hcl oral*
- *pentazocine-naloxone hcl*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Mild pain: codeine. Moderate to severe pain: short-term NSAIDs, tramadol, tramadol/APAP, morphine sulfate, hydrocodone/APAP, oxycodone, oxycodone/APAP, fentanyl.

HRM-ANTI-ARRHYTHMICS

Products Affected

- DIGITEK ORAL TABLET 250 MCG
- DIGOX ORAL TABLET 250 MCG
- *digoxin oral solution*
- *digoxin oral tablet 250 mcg*
- *disopyramide phosphate oral*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-ANTIEMETIC DRUGS

Products Affected

- *promethazine hcl oral syrup*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-ANTI-HISTAMINES

Products Affected

- *carbinoxamine maleate oral solution*
- *clemastine fumarate oral tablet 2.68 mg*
- *cyproheptadine hcl oral solution*
- *promethazine vc plain oral solution*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine

HRM-ANTIHYPERTENSIVE AGENTS

Products Affected

- *methyldopa oral*
- *methyldopa-hydrochlorothiazide*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-ANTIPARKINSON AGENTS

Products Affected

- *benztropine mesylate oral*
- *trihexyphenidyl hcl*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For the treatment of Parkinsonism, a trial of at least 1 formulary drug that is not an HRM in elderly (e.g., carbidopa/levodopa, pramipexole, ropinirole, bromocriptine, selegiline, rasagiline, entacapone, amantadine, etc.) or other type of clinical justification will be required in members 65 years of age and older. For the treatment of medication-induced movement disorder - extrapyramidal disease, a trial of at least 1 formulary drug that is not an HRM in elderly (e.g., amantadine, etc.) or other type of clinical justification will be required in members 65 years of age and older.

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HRM-CALCIUM CHANNEL BLOCKERS, DIHYDROPYRIDINE

Products Affected

- *nifedipine oral*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	extended-release nifedipine, nicardipine, amlodipine

HRM-MEGESTROL

Products Affected

- *megestrol acetate oral suspension 625 mg/5ml*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND If formulary non HRM alternatives for diagnosis of cachexia secondary to chronic illness (dronabinol, oxandrolone) considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 of the alternative(s) AND the prescribing physician attests to the medical necessity for using this high risk medication, AND anticipated treatment course/duration. For treatment of cancer related diagnosis or endometrial hyperplasia, or endometriosis, requests will be automatically approved.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

HRM-ORAL AND TRANSDERMAL ESTROGENS AND PROGESTINS

Products Affected

- CLIMARA PRO
- *estradiol oral*
- *estropipate oral tablet 0.75 mg*
- MENEST ORAL TABLET 0.3 MG, 0.625 MG, 1.25 MG
- PREMARIN ORAL
- PREMPRO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Bone Density: alendronate, risedronate, ibandronate, raloxifene (zoledronic acid for bed-bound patients or for post-hip fracture). Vaginal Symptoms: vaginal estrogen cream

HRM-PLATELET INHIBITORS

Products Affected

- *dipyridamole oral*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	clopidogrel, aggrenox

HRM-SEDATIVE HYPNOTIC AGENTS

Products Affected

- *zaleplon*
- *zolpidem tartrate er*
- *zolpidem tartrate oral*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND If formulary non HRM alternatives, Silenor (less than or equal to 6mg/d) considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to the alternative(s) AND the prescribing physician attests to the medical necessity for using this high risk medication, AND Monitoring plan for adverse side effects, AND anticipated treatment course/duration.
Age Restrictions	PA applies to patients 65 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Non-HRM alternatives, must inadequate response, intolerable side effect, or contraindication to both: Rozerem (8 mg/d), Silenor (less than or equal to 6mg/d)

HRM-SKELETAL MUSCLE RELAXANTS

Products Affected

- *carisoprodol oral*
- *carisoprodol-aspirin*
- *carisoprodol-aspirin-codeine*
- *chlorzoxazone oral tablet 500 mg*
- *cyclobenzaprine hcl oral*
- *methocarbamol oral*
- *orphenadrine citrate er*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-SULFONYLUREAS

Products Affected

- *chlorpropamide*
- *glyburide oral*
- *glyburide micronized*
- *glyburide-metformin*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	glimepiride, glipizide

HRM-UTI ANTIBACTERIALS

Products Affected

- *nitrofurantoin macrocrystal oral*
- *nitrofurantoin monohyd macro*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND If formulary non HRM alternatives (Non-HRM alternatives: sulfamethoxazole/trimethoprim) considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to the alternative AND the prescribing physician attests to the medical necessity for using this high risk medication and intent to monitor for side effects, AND anticipated treatment course/duration
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

IBRANCE

Products Affected

- IBRANCE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND One of the following: 1) Used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy OR 2) Used in combination with an aromatase inhibitor AND One of the following: 1) patient is a postmenopausal woman OR 2) both of the following: patient is a premenopausal or perimenopausal woman and patient is receiving a luteinizing hormone-releasing hormone (LHRH) agonist [eg, Zoladex (goserelin)].
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ICLUSIG

Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic myelogenous leukemia (CML) AND One of the following: A) History of failure, resistance, relapse, contraindication, or intolerance to at least TWO other tyrosine kinase inhibitors (SPRYCEL, TASIGNA, and BOSULIF), or B) Patient has the T315I mutation. OR Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) AND One of the following: A) History of failure, resistance relapse, contraindication, or intolerance to at least TWO other FDA-approved tyrosine kinase inhibitors (SPRYCEL, TASIGNA, BOSULIF), or B) Patient has the T315I mutation.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

IDHIFA

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia with an isocitrate dehydrogenase 2 mutation as detected by an FDA approved test.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

IMBRUVICA

Products Affected

- IMBRUVICA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of mantle cell lymphoma (MCL) who have received at least one prior therapy OR chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) OR chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion OR Waldenstrom's macroglobulinemia (WM) OR marginal zone lymphoma who require systemic therapy and have received at least one prior anti-CD20-based therapy OR graft vs host disease after failure of a least one first-line corticosteroid therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

INLYTA

Products Affected

- INLYTA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced renal cell carcinoma AND patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

INTRAROSA

Products Affected

- INTRAROSA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, known or suspected estrogen-dependent neoplasia.
Required Medical Information	Diagnosis of moderate to severe dyspareunia or atrophic vaginitis AND A) Patient must be female, B) Patient must be menopausal or postmenopausal, C) Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (e.g. Premarin vaginal cream, Estrace vaginal cream, Estring, Vagifem), D) Patient does not have renal or hepatic impairment.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Reauthorization: 12 months
Other Criteria	None

INVANZ

Products Affected

- INVANZ INJECTION

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Invanz (ertapenem sodium) is contraindicated in patients with a hypersensitivity to ertapenem or other drugs of the same class, in patients with a prior anaphylactic reaction to beta-lactams and in patients with a hypersensitivity to amide-type anesthetics.
Required Medical Information	Labs with culture and sensitivity information.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

IRESSA

Products Affected

- IRESSA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

JAKAFI

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Myelofibrosis: Diagnosis of primary myelofibrosis, OR post-polycythemia vera myelofibrosis, OR post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera, AND history of failure, contraindication, or intolerance to hydroxyurea.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	This criteria applies to new starts only

JUXTAPID

Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 5 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests. Pregnancy. Concomitant use with strong or moderate CYP3A4 inhibitors.
Required Medical Information	Diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (LDLRAP1 or ARH), or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient has been receiving at least 12 consecutive weeks of ONE LDL-C lowering prescription therapy and will continue to receive an LDL-C lowering prescription therapy. History of failure after 12 consecutive weeks or intolerance to PCSK9 inhibitor therapy. One of the following LDL-C values while on maximally tolerated lipid-lowering regimen within the last 30 days: LDL-C greater than or equal to 100 mg/dL with ASCVD OR LDL-C greater than or equal to 130 mg/dL without ASCVD. Not used in combination with Kynamro (mipomersen). Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. Patient is not pregnant. Patient does not have moderate or severe hepatic impairment (Child-Pugh category B or C) or active liver disease including unexplained persistent abnormal liver function tests. Patient is not concomitantly on moderate or strong CYP 3A4 inhibitors (ie. clarithromycin).

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	None

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis AND the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data.
Age Restrictions	Oral granules: 2 years of age and older. Oral tablets: 6 years of age and older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

KISQALI

Products Affected

- KISQALI 200 DOSE
- KISQALI 400 DOSE
- KISQALI 600 DOSE
- KISQALI FEMARA 200 DOSE
- KISQALI FEMARA 400 DOSE
- KISQALI FEMARA 600 DOSE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or advanced, HER-2 negative, hormone receptor-positive, postmenopausal women in combination with fulvestrant as initial endocrine based therapy or following disease progression on endocrine therapy (requirement of fulvestrant applies to single agent Kisqali only, NOT Kisqali-Femara Co-Pack) OR Diagnosis metastatic or advanced, HER-2 negative, hormone receptor-positive, premenopausal, perimenopausal, or postmenopausal women, in combination with an aromatase inhibitor for initial endocrine-based treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or gastroenterologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

KORLYM

Products Affected

- KORLYM

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Supporting statement of diagnosis and relevant medical information from physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

KYNAMRO

Products Affected

- KYNAMRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests.
Required Medical Information	Diagnosis of homozygous familial hypercholesterolemia as evidenced by one of the following: A) genetic confirmation of 2 mutant alleles at the LDL receptor, ApoB, PCSK9, or autosomal recessive hypercholesterolemia (ARH) adaptor protein gene locus OR B) untreated/pre-treatment LDL greater than 500 mg/dL with at least one of the following: cutaneous or tendonous xanthoma before age 10 years, history of early vascular disease (men younger than 55 years, women younger than 60 years) on both sides of the family if parenteral LDL levels are unknown, elevated LDL cholesterol levels before lipid-lowering therapy consistent with heterozygous FH in both parents AND Patient has tried and had an inadequate response to the maximum tolerated dose of a high potency statin (e.g., atorvastatin, rosuvastatin), unless all statins are contraindicated.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Other Criteria	For renewal, patient has responded to therapy with a decrease in LDL levels from baseline AND patient does not have contraindications to therapy.

LENVIMA

Products Affected

- LENVIMA 10 MG DAILY DOSE
- LENVIMA 14 MG DAILY DOSE
- LENVIMA 18 MG DAILY DOSE
- LENVIMA 20 MG DAILY DOSE
- LENVIMA 24 MG DAILY DOSE
- LENVIMA 8 MG DAILY DOSE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer OR advanced renal cell carcinoma following one prior anti-angiogenic therapy in combination with everolimus
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

LETAIRIS

Products Affected

- LETAIRIS

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	This criteria applies to new starts only

LIDOCAINE PATCH

Products Affected

- *lidocaine external patch 5 %*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pain associated with diabetic neuropathy OR pain associated with cancer-related neuropathy OR post-herpetic neuralgia.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

LONSURF

Products Affected

- LONSURF

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For initial treatment: Absolute neutrophil count 1,500/mm ³ or greater or febrile neutropenia resolved, platelet count 75,000/mm ³ or greater, and grade 3 or 4 nonhematological reactions resolved to grade 0 or 1

LUPRON

Products Affected

- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.
Required Medical Information	Diagnosis of one of the following: A) advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), B) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only) and for initial, patient has had an inadequate pain control response or patient has an intolerance or contraindication to one of the following: Danazol OR Combination [estrogen/progesterone] Oral Contraceptives OR Progestins and for retreatment course, Patient is experiencing recurrence of symptoms after an initial course of therapy with leuprolide acetate and Norethindrone acetate 5 mg daily will be co-administered, or C) Anemia due to uterine Leiomyomata (Fibroids) (3.75 mg 1-month & 11.25 mg 3-month depots only) and Patient is preoperative
Age Restrictions	Uterine fibroids, endometriosis and prostate cancer: 18 years of age and older, CPP: age 2-11 female and 2-12 male
Prescriber Restrictions	None
Coverage Duration	Endometriosis- 6 months, Uterine fibroids -3 months, Prostate cancer, Precocious Puberty -12 months

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Other Criteria	For endometriosis and uterine fibroids, patient will be using nonhormonal contraception during and for 12 weeks after therapy.

LYNPARZA

Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Appropriate diagnosis and testing for BRCA mutation (deleterious or suspected deleterious germline BRCA mutated (as detected by an FDA approved test) advanced ovarian cancer that has been treated with 3 or more prior lines of chemotherapy)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

MATULANE

Products Affected

- MATULANE

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Treatment of Hodgkin's Lymphoma OR medulloblastoma in combination with nitrogen mustard, vincristine and prednisone OR high-grade malignant glioma in combination with lomustine and vincristine
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

MEKINIST

Products Affected

- MEKINIST

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic anaplastic thyroid cancer with documented positive BRAF V600E or V600K mutation as detected by an FDA-approved test as single agent or used in combination with Tafinlar OR Diagnosis of metastatic or unresectable melanoma and metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation as detected by an FDA-approved test OR Diagnosis of BRAF V600K mutation-positive unresectable or metastatic melanoma or use as adjuvant treatment of BRAF V600K mutation-positive melanoma
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

MS INTERFERONS

Products Affected

- AVONEX
- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT
- PLEGRIDY
- PLEGRIDY STARTER PACK SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NERLYNX

Products Affected

- NERLYNX

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of early stage HER2- overexpressed breast cancer. Must be used after trastuzumab therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

NEXAVAR

Products Affected

- NEXAVAR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel.
Required Medical Information	Diagnosis of unresectable hepatocellular carcinoma OR Diagnosis of advanced renal cell carcinoma OR Diagnosis of locally recurrent or metastatic, progressive, differentiated thyroid carcinoma refractory to radioactive iodine treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

NINLARO

Products Affected

- NINLARO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, documentation of combination therapy with lenalidomide and dexamethasone. History of 1 prior therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

NORTHERA

Products Affected

- NORTHERA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Request will be approved for the following indication(s): orthostatic dizziness, light-headedness, or the feeling that you are about to black out in adults with neurogenic orthostatic hypotension (NOH) caused by primary autonomic failure (i.e., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, and non-diabetic autonomic neuropathy).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient diagnosis of pseudobulbar affect.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NUPLAZID

Products Affected

- NUPLAZID ORAL TABLET 17 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Parkinson disease psychosis including hallucinations and/or delusions
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

OPSUMIT

Products Affected

- OPSUMIT

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy.
Required Medical Information	Diagnosis of pulmonary arterial hypertension WHO group I AND diagnosis was confirmed by right heart catheterization AND female patients are enrolled in the OPSUMIT REMS program.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	None

ORKAMBI

Products Affected

- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Initial Therapy: Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test AND if less than 18 years of age, baseline ophthalmological exam completed. Continuation of therapy: Documentation patient is tolerating and responding to medication (i.e. improved FEV1, weight gain, decreased exacerbations, etc.)
Age Restrictions	None
Prescriber Restrictions	Must be prescribed by, or in conjunction with, a pulmonologist or is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None

OSPHERA

Products Affected

- OSPHERA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, known or suspected estrogen-dependent neoplasia, acute thromboembolism or a past history of thromboembolic disease (including patients with a history of DVT, pulmonary embolism, retinal vein thrombosis, stroke, or myocardial infarction, known or suspected pregnancy.
Required Medical Information	Diagnosis of moderate to severe dyspareunia or atrophic vaginitis AND A) Patient must be female, B) Patient must be menopausal or postmenopausal, C) Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (Premarin vaginal cream, Estrace vaginal cream, Estring, Vagifem), D) Dose must not exceed 1 tablet per day, E) Patient does not have hepatic impairment.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

PCSK9 INHIBITOR

Products Affected

- PRALUENT SUBCUTANEOUS SOLUTION PEN-INJECTOR
- REPATHA
- REPATHA PUSHTRONEX SYSTEM
- REPATHA SURECLICK

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>For PRALUENT: MUST MEET CRITERIA #1 OR #3. For REPATHA: MUST MEET CRITERIA #1, #2, OR #3. 1. Diagnosis of heterozygous familial hypercholesterolemia (HeFH) confirmed by genotyping OR Simon Broome criteria: Total cholesterol greater than 290mg/dL or LDL cholesterol greater than 190mg/dL, PLUS ONE OF THE FOLLOWING: Tendon xanthomas in patient or 1st degree relative (parent, sibling, child) or 2nd degree relative (grandparent, uncle, aunt) OR DNA-based evidence of LDL receptor mutation, familial defective apo B-100, or PCSK9 mutation. 2a. Myocardial infarction prophylaxis, stroke prophylaxis, and to reduce risk of coronary revascularization in patients with established CVD OR 2b. Diagnosis of homozygous familial hypercholesterolemia (HoFH) confirmed by genotyping OR diagnosis based on the following: a. History of untreated LDL-C greater than 500 mg/dL AND xanthoma before 10 years of age OR b. Documentation of HeFH in both parents. 3. Diagnosis of clinical atherosclerotic cardiovascular disease as defined as one of the following: a. acute coronary syndrome, b. history of myocardial infarction, c. stable/unstable angina, d. coronary or other arterial revascularization, e. stroke, f. transient ischemic stroke, g. peripheral arterial disease presumed to be atherosclerotic region AND MEETS CRITERIA #4, #5, and #6. 4. Provide baseline and current LDL-C. 5. LDL-C greater than or equal to 70mg/dL. 6. Used</p>

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
	in combination with maximally tolerated high-intensity statin OR MEETS CRITERIA #7 AND #8. 7. Statin intolerant 8. LDL-C greater than or equal to 70mg/dL. CONTINUING THERAPY: 1. Documented response to Praluent or Repatha, defined as ONE of the following: a. The patient is tolerating medication b. Will continue to be used in combination with maximally tolerated statin (unless statin intolerant).
Age Restrictions	Repatha: 13 years of age and older for diagnosis HoFM, Diagnosis CVD and HeFH AND Praluent and Repatha : 18 years of age and older
Prescriber Restrictions	Must be prescribed by, or in consultation with, a cardiologist, endocrinologist, or lipid specialist
Coverage Duration	Initial approval: 8 weeks, Renewal approval: 12 months
Other Criteria	None

POMALYST

Products Affected

- POMALYST

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Documentation of ALL of the following: 1. Disease has progressed within 60 days of completion of the last therapy 2. If female of reproductive potential ALL of the below: Two negative pregnancy tests obtained prior to initiating therapy with Pomalyst, monthly negative pregnancy tests during therapy 3. Patient has been counseled about the use of reliable contraception before, during, and 1 month after initiation of therapy with Pomalyst 4. Patient assessment to determine if prophylactic aspirin or antithrombic treatment (warfarin, clopidogrel) will need to be taken to reduce the risk of VTE (embolism, stroke) 5. Registered and certified to be compliant with Pomalyst REMS (Risk Evaluation and Mitigation Strategy) program
Age Restrictions	None
Prescriber Restrictions	Oncologist or hematologist
Coverage Duration	12 months
Other Criteria	A documented diagnosis of multiple myeloma and received at least two prior therapies including lenalidomide (Revlimid) and bortezomib (Velcade)

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

PROVIGIL

Products Affected

- *modafinil*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

REGANEX

Products Affected

- REGANEX

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diabetic Neuropathic Ulcers: Diabetic patient with ulcer wound. Treatment will be given in combination with ulcer wound care (debridement, infection control, and/or pressure relief).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Diabetic Neuropathic Ulcers: Maximum 5 months.
Other Criteria	None

RUBRACA

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious BRCA mutation (germline and/or somatic)-associated epithelial ovarian, fallopian tube, or primary peritoneal cancer and all of the following criteria: 1. BRCA mutation detected by an approved FDA laboratory test, 2. Previous trial/failure with two or more chemotherapy regimens, 3. Used as monotherapy, 4. Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, 5. Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose OR Diagnosis of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer and all of the following: 1. Complete or partial response to platinum-based chemotherapy 2. Used as monotherapy 3. Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, 4. Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Hematologist or Oncologist
Coverage Duration	12 months

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Other Criteria	This criteria applies to new starts only

RYDAPT

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Angioedema
Required Medical Information	Diagnosis of treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and consolidation therapy or diagnosis of systemic mastocytosis.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

SAMSCA

Products Affected

- SAMSCA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SILDENAFIL

Products Affected

- *sildenafil citrate oral tablet 20 mg*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SPRYCEL

Products Affected

- SPRYCEL

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) that is newly diagnosed in the chronic phase OR Ph+ CML with resistance or intolerance to prior therapy, including imatinib OR Diagnosis of Ph+ acute lymphoblastic leukemia with resistance or intolerance to prior therapy OR Gastrointestinal stromal tumors (GIST) after disease progression on imatinib or Sutent (sunitinib)
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

STIVARGA

Products Affected

- STIVARGA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer AND documentation of prior therapy with ALL of the following per the indication: 1. fluoropyrimidine, oxaliplatin, and irinotecan-based chemotherapy 2. bevacizumab (Avastin) 3. panitumumab (Vectibix) OR cetuximab (Erbix) (for KRAS mutation-negative patients only) OR a documented diagnosis of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) who have been previously treated with imatinib mesylate (Gleevec) and sunitinib malate (Sutent) OR a documented diagnosis of hepatocellular carcinoma in patients previously treated with sorafenib (Nexavar).
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

SUTENT

Products Affected

- SUTENT

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of gastrointestinal stromal tumors after disease progression on or intolerance to Gleevec OR Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease OR Diagnosis of high risk recurrent renal cell carcinoma following nephrectomy, used as adjuvant therapy.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

SYLATRON

Products Affected

- SYLATRON SUBCUTANEOUS KIT
200 MCG, 300 MCG, 600 MCG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Autoimmune hepatitis. Hepatic decompensation (Child-Pugh score greater than 6 [Class B or C])
Required Medical Information	Melanoma: Diagnosis of melanoma with microscopic or gross nodal involvement AND The prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and patient is homozygous for the F508del mutation OR have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	None

SYNRIBO

Products Affected

- SYNRIPO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic myelogenous leukemia AND patient has tried and failed or has a contraindication or intolerance to 2 tyrosine kinase inhibitors
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

SYPRINE

Products Affected

- *trientine hcl*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease and intolerance to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TAFINLAR

Products Affected

- TAFINLAR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic anaplastic thyroid cancer with documented positive BRAF V600E or V600K mutation as detected by an FDA-approved test as single agent or used in combination with Mekinist OR Diagnosis of metastatic or unresectable melanoma and metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation as detected by an FDA-approved test
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	Tafinlar should not be used in patients with wild-type BRAF melanoma due to the potential risk of tumor promotion in these patients

TAGRISSO

Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, non-small cell lung cancer with one of the following- confirmed presence of T790M EGFR tumor mutation OR confirmed presence of epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R tumor mutations, as detected by an FDA-approved test.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

TARCEVA

Products Affected

- TARCEVA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer and Tarceva will be used in combination with gemcitabine OR Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer with one of the following: A) failure with at least one prior chemotherapy regimen and Tarceva will be used as monotherapy, or B) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and Tarceva will be used as maintenance treatment and Tarceva will be used as monotherapy, or C) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

TASIGNA

Products Affected

- TASIGNA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Long QT syndrome. Uncorrected hypokalemia. Uncorrected hypomagnesemia. Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors
Required Medical Information	Diagnosis of newly diagnosed Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) in the chronic phase OR Diagnosis of Ph+ CML with resistance or intolerance to prior therapy that include imatinib.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

TECFIDERA

Products Affected

- TECFIDERA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondary-progressive MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TOBI

Products Affected

- TOBI PODHALER

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	This agent may be considered medically necessary when the following criteria are met: Cystic Fibrosis-The patient has a diagnosis of cystic fibrosis (CF). The patient is colonized with P. Aeruginosa.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TRACLEER

Products Affected

- TRACLEER

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Receiving concomitant cyclosporine A or glyburide therapy. Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal.
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class II-IV AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	Liver aminotransferases will be measured prior to initiation of treatment and then monthly.

TRETINOIN

Products Affected

- *tretinoin external cream*
- *tretinoin external gel 0.01 %, 0.025 %*

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Approval will be given to all members using this agent for a medically necessary, FDA approved, non-cosmetic indication(s).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TYKERB

Products Affected

- TYKERB

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND a) the medication will be used in combination with Xeloda in a patient with advanced or metastatic disease and the patient has received prior therapy including an anthracycline, a taxane, and trastuzumab or b) The medication will be used in combination with Femara for the treatment of a postmenopausal woman with hormone receptor-positive metastatic disease for whom hormonal therapy is indicated.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

TYMLOS

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients at increased risk of osteogenic sarcoma.
Required Medical Information	Diagnosis of osteoporosis in post-menopausal women at high risk for fracture. Member must have failed therapy with a bisphosphonate (defined by a fracture while on therapy or worsening bone density) unless such a trial is shown to be inappropriate or contraindicated (i.e., presence of severe osteoporosis [T-scores -3.0 or worse in lumbar spine, femoral neck, or total hip region], history of major osteoporotic fracture, presence of renal insufficiency, etc) AND member has at least one of the following: T-score equal to or worse than -2.5 in the lumbar spine, femoral neck, or total hip region OR a FRAX calculator based 10-year risk of at least 20% for a major osteoporotic fracture (spine, shoulder, hip, or wrist), or a 10-year risk of at least 3% for a hip fracture OR presence or history of osteoporotic fracture.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: Total duration not to exceed 24 months during patient's lifetime.
Other Criteria	None

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

UPTRAVI

Products Affected

- UPTRAVI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization AND patient has tried and had an insufficient response to at least one other PAH agent (e.g., sildenafil) therapy and one other ERA agent (e.g. letairis, opsumit, tracleer).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

VENCLEXTA

Products Affected

- VENCLEXTA
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic lymphocytic leukemia (CLL) OR small lymphocytic lymphoma, with or without 17p deletion and patient has had at least 1 prior therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

VERZENIO

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	BREAST CANCER (1) Patient must have a diagnosis of advanced or metastatic breast cancer AND (2a) must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy OR (2b) used as monotherapy for treatment of disease progression following endocrine therapy and patient has already received at least one prior chemotherapy regimen of Ibrance or Kisqali OR (2c) used as initial endocrine-based treatment in combination with an aromatase inhibitor AND (3) disease is hormone receptor positive AND human epidermal growth factor 2 (HER2)- negative
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of therapy

VOTRIENT

Products Affected

- VOTRIENT

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (doxorubicin, dacarbazine, ifosfamide, epirubicin, docetaxel, or vinorelbine).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

XALKORI

Products Affected

- XALKORI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer AND patient has non-squamous cell histology AND Patient has anaplastic lymphoma kinase (ALK)-positive disease as detected with an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility or are ROS1-positive
Age Restrictions	18 years of age and older
Prescriber Restrictions	Must be prescribed by an oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

XTANDI

Products Affected

- XTANDI

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Metastatic castration-resistant prostate cancer (mCRPC): Diagnosis of mCRPC. History of failure, contraindication or intolerance to Zytiga.
Age Restrictions	None
Prescriber Restrictions	Prescribed or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

XURIDEN

Products Affected

- XURIDEN

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Hereditary orotic aciduria
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a specialist that treats metabolic defects
Coverage Duration	12 months
Other Criteria	None

YONSA

Products Affected

- YONSA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic castration-resistant prostate cancer, and used in combination with methylprednisolone AND Documented history of trial with, inadequate treatment response, adverse event, or contraindication to Zytiga
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ZEJULA

Products Affected

- ZEJULA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of recurrent epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer and patient had a complete or partial response to platinum-based chemotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or gynecologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ZELBORAF

Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic melanoma OR Erdheim-Chester disease. Patient has positive BRAF-V600E mutation documented by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ZOLINZA

Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma AND progressive, persistent or recurrent disease or patient is not a candidate for or following 2 systemic therapies (bexarotene, romidepsin, etc.)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ZYDELIG

Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The patient has one of the following diagnoses: A) chronic lymphocytic leukemia AND The medication will be used in combination with rituximab AND The patient has relapsed on at least one prior therapy (purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]) AND the patient does not have any co-morbidities that prevents the use of cytotoxic chemotherapy (severe neutropenia or thrombocytopenia, creatinine clearance less than 60 mL/minute), B) follicular lymphoma AND the patient has relapsed on at least two prior systemic therapies (rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]), or C) small lymphocytic lymphoma AND The patient has relapsed on at least two prior systemic therapies (rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months

Formularies: ID 19482 V.8, ID 19485 V.8, ID 19487 V.8.
Last Updated: 12/12/2018
Version: 8 - Effective: 01/01/2019

HealthSun Health Plans
2019 Prior Authorization Criteria

PA Criteria	Criteria Details
Other Criteria	This criteria applies to new starts only

ZYKADIA

Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

ZYTIGA

Products Affected

- ZYTIGA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of metastatic prostate cancer AND Patient has castration-resistant disease (defined by tumor growth/disease progression, risk in PSA levels, new metastases) OR high-risk castration-sensitive prostate cancer AND Zytiga will be used in combination with prednisone.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only

PART B VERSUS PART D

Products Affected

- *acetylcysteine inhalation*
- *acyclovir sodium intravenous solution*
- *albuterol sulfate inhalation*
- AMBISOME
- *amikacin sulfate injection solution 500 mg/2ml*
- AMINOSYN II INTRAVENOUS SOLUTION 10 %, 8.5 %
- AMINOSYN II/ELECTROLYTES
- AMINOSYN/ELECTROLYTES
- AMINOSYN-HBC
- AMINOSYN-PF
- AMINOSYN-RF
- *amphotericin b injection*
- *ampicillin sodium injection solution reconstituted 1 gm, 125 mg*
- *ampicillin sodium intravenous solution reconstituted 10 gm*
- *ampicillin-sulbactam sodium injection*
- *aprepitant*
- ARCALYST
- ASTAGRAF XL
- AZACTAM INJECTION SOLUTION RECONSTITUTED 2 GM
- AZASAN
- *azathioprine oral*
- *azithromycin intravenous solution reconstituted 500 mg*
- BIVIGAM INTRAVENOUS SOLUTION 10 GM/100ML
- BROVANA
- *budesonide inhalation*
- *calcitonin (salmon)*
- *calcitriol oral*
- *casprofungin acetate*
- *cefazolin sodium injection solution reconstituted 1 gm, 10 gm, 500 mg*
- *cefepime hcl injection*
- *cefoxitin sodium*
- *ceftriaxone sodium injection solution reconstituted 1 gm, 2 gm, 250 mg, 500 mg*
- *ceftriaxone sodium intravenous solution reconstituted 10 gm*
- *cefuroxime sodium injection solution reconstituted 7.5 gm, 750 mg*
- *cefuroxime sodium intravenous solution reconstituted 1.5 gm*
- *chlorpromazine hcl oral*
- CINRYZE
- *ciprofloxacin in d5w intravenous solution 200 mg/100ml*
- *clindamycin phosphate injection solution 600 mg/4ml*
- CLINIMIX E/DEXTROSE (2.75/10)
- CLINIMIX E/DEXTROSE (2.75/5)
- CLINIMIX E/DEXTROSE (4.25/10)
- CLINIMIX E/DEXTROSE (4.25/25)
- CLINIMIX E/DEXTROSE (4.25/5)
- CLINIMIX E/DEXTROSE (5/15)
- CLINIMIX E/DEXTROSE (5/20)
- CLINIMIX E/DEXTROSE (5/25)
- CLINIMIX/DEXTROSE (2.75/5)
- CLINIMIX/DEXTROSE (4.25/10)
- CLINIMIX/DEXTROSE (4.25/20)
- CLINIMIX/DEXTROSE (4.25/25)
- CLINIMIX/DEXTROSE (4.25/5)
- CLINIMIX/DEXTROSE (5/15)
- CLINIMIX/DEXTROSE (5/20)
- CLINIMIX/DEXTROSE (5/25)
- CLINISOL SF
- *colistimethate sodium (cba)*
- *cromolyn sodium inhalation*

HealthSun Health Plans
2019 Prior Authorization Criteria

- *cyclophosphamide oral capsule*
- *cyclosporine modified*
- *cyclosporine oral capsule*
- *daptomycin intravenous solution reconstituted 500 mg*
- DEPO-PROVERA INTRAMUSCULAR SUSPENSION 400 MG/ML
- *dextrose intravenous solution 10 %, 5 %*
- *dextrose-nacl intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225 %, 5-0.33 %, 5-0.45 %, 5-0.9 %*
- *diphtheria-tetanus toxoids dt*
- *doripenem intravenous solution reconstituted 500 mg*
- *doxercalciferol oral*
- DOXY 100
- *dronabinol*
- *duramorph*
- ELIGARD
- ENGERIX-B INJECTION SUSPENSION 10 MCG/0.5ML, 20 MCG/ML
- ENVARUS XR
- ERAXIS
- ERYTHROCIN LACTOBIONATE INTRAVENOUS SOLUTION RECONSTITUTED 500 MG
- FIRMAGON
- FLEBOGAMMA DIF INTRAVENOUS SOLUTION 5 GM/50ML
- *fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%*
- FREAMINE HBC
- *furosemide injection*
- GAMMAGARD INJECTION SOLUTION 2.5 GM/25ML
- GAMMAGARD S/D LESS IGA
- GAMMAKED INJECTION SOLUTION 1 GM/10ML
- GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML
- GAMUNEX-C INJECTION SOLUTION 1 GM/10ML
- GENGRAF ORAL CAPSULE 100 MG, 25 MG
- GENGRAF ORAL SOLUTION
- *gentamicin sulfate injection solution 40 mg/ml*
- *granisetron hcl oral*
- *heparin sodium (porcine) injection solution 1000 unit/ml, 10000 unit/ml, 20000 unit/ml, 5000 unit/ml*
- HEPATAMINE
- *imipenem-cilastatin intravenous solution reconstituted 250 mg*
- IMOVAX RABIES
- INTRON A
- IONOSOL-MB IN D5W
- *ipratropium bromide inhalation*
- *ipratropium-albuterol*
- ISOLYTE-P IN D5W
- ISOLYTE-S
- *kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.33 meq/l-%-%, 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%*
- *kcl-lactated ringers-d5w*
- LEUKINE INTRAVENOUS
- *leuprolide acetate injection*
- *levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/0.5ml, 1.25 mg/3ml*
- *levocarnitine oral solution*
- *levocarnitine oral tablet*

HealthSun Health Plans
2019 Prior Authorization Criteria

- *levofloxacin in d5w intravenous solution 500 mg/100ml, 750 mg/150ml*
- *levofloxacin intravenous*
- *linezolid intravenous solution 600 mg/300ml*
- *magnesium sulfate injection solution 50 %, 50 % (10ml syringe)*
- *meropenem*
- *methotrexate oral*
- *methotrexate sodium (pf) injection solution 50 mg/2ml*
- *methotrexate sodium injection solution 250 mg/10ml*
- *metronidazole in nacl intravenous solution 500-0.79 mg/100ml-%*
- MIRCERA INJECTION SOLUTION PREFILLED SYRINGE 100 MCG/0.3ML
- *moxifloxacin hcl in nacl*
- MYCAMINE
- *mycophenolate mofetil*
- *mycophenolate sodium*
- *nafcillin sodium injection solution reconstituted 1 gm*
- *nafcillin sodium intravenous solution reconstituted 10 gm*
- NEBUPENT
- NEPHRAMINE
- NORMOSOL-M IN D5W
- NORMOSOL-R IN D5W
- NORMOSOL-R PH 7.4
- *nutrilipid intravenous emulsion 20 %*
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML
- *octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml*
- *ondansetron*
- *ondansetron hcl oral*
- *paricalcitol oral*
- *penicillin g potassium injection solution reconstituted 20000000 unit*
- *penicillin g sodium*
- PENTAM
- *perphenazine oral tablet 4 mg, 8 mg*
- *piperacillin sod-tazobactam so intravenous solution reconstituted 2.25 (2-0.25) gm, 3.375 (3-0.375) gm, 4.5 (4-0.5) gm*
- PLASMA-LYTE 148
- PLASMA-LYTE A
- PLENAMINE
- *potassium chloride in dextrose intravenous solution 20-5 meq/l-%, 40-5 meq/l-%*
- *potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%*
- *potassium chloride intravenous solution 2 meq/ml, 20 meq/100ml*
- PREMASOL
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML
- PROCALAMINE
- *prochlorperazine maleate oral*
- PROLASTIN-C INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG
- PROSOL
- PULMOZYME
- RABAVERT
- RAPAMUNE ORAL SOLUTION
- RECOMBIVAX HB
- *rifampin intravenous*
- SENSIPAR
- *sirolimus oral*
- *sodium chloride injection solution 2.5 meq/ml*

HealthSun Health Plans
2019 Prior Authorization Criteria

- *sodium chloride intravenous solution 0.45 %, 0.9 %, 3 %, 5 %*
- *sodium lactate intravenous solution 5 meq/ml*
- SYNDROS
- *tacrolimus oral*
- TEFLARO
- TENIVAC
- *tetanus-diphtheria toxoids td*
- *tigecycline*
- *tobramycin inhalation*
- *tobramycin sulfate injection solution 10 mg/ml, 80 mg/2ml*
- TPN ELECTROLYTES INTRAVENOUS SOLUTION
- TRAVASOL
- TRELSTAR MIXJECT
- TREXALL
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- TWINRIX
- *vancomycin hcl intravenous solution reconstituted 10 gm, 1000 mg, 500 mg*
- VARUBI ORAL
- XATMEP
- ZORTRESS ORAL TABLET 0.25 MG, 0.5 MG, 0.75 MG

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index

A

acetylcysteine inhalation	130
acyclovir sodium intravenous solution.....	130
adefovir dipivoxil.....	34
ADEMPAS.....	2
AFINITOR.....	3, 4
AFINITOR DISPERZ	3, 4
albuterol sulfate inhalation	130
ALECENSA	5
ALUNBRIG.....	6
AMBISOME.....	130
amikacin sulfate injection solution 500 mg/2ml	130
AMINOSYN II INTRAVENOUS SOLUTION 10 %, 8.5 %	130
AMINOSYN II/ELECTROLYTES ..	130
AMINOSYN/ELECTROLYTES.....	130
AMINOSYN-HBC.....	130
AMINOSYN-PF	130
AMINOSYN-RF	130
amphotericin b injection.....	130
ampicillin sodium injection solution reconstituted 1 gm, 125 mg..	130
ampicillin sodium intravenous solution reconstituted 10 gm.	130
ampicillin-sulbactam sodium injection.....	130
AMPYRA.....	7
aprepitant.....	130
ARCALYST.....	130
ASCOMP-CODEINE	38
ASTAGRAF XL.....	130
AUBAGIO	8
AURYXIA.....	9
AVONEX.....	78
AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT	78
AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT	78

AZACTAM INJECTION SOLUTION

RECONSTITUTED 2 GM	130
AZASAN	130
azathioprine oral	130
azithromycin intravenous solution reconstituted 500 mg	130

B

benztropine mesylate oral.....	43
BETASERON SUBCUTANEOUS KIT	78
bexarotene	10
BIVIGAM INTRAVENOUS SOLUTION 10 GM/100ML.....	130
BOSULIF	11
BRIVIACT ORAL.....	12
BROVANA.....	130
budesonide inhalation	130
butalbital-acetaminophen oral tablet 50-325 mg.....	38
butalbital-apap-caff-cod	38
butalbital-apap-caffeine oral capsule	38
butalbital-apap-caffeine oral tablet 50-325-40 mg	38
butalbital-asa-caff-codeine.....	38

C

CABOMETYX	13
calcitonin (salmon)	130
calcitriol oral	130
CALQUENCE.....	14
CAPRELSA	15
carbinoxamine maleate oral solution	41
CARIMUNE NF INTRAVENOUS SOLUTION RECONSTITUTED 6 GM	16, 17
carisoprodol oral.....	50
carisoprodol-aspirin	50
carisoprodol-aspirin-codeine.....	50
caspofungin acetate	130

HealthSun Health Plans
2019 Prior Authorization Criteria

cefazolin sodium injection solution reconstituted 1 gm, 10 gm, 500 mg	130	CLINIMIX/DEXTROSE (4.25/5)	130
cefepime hcl injection	130	CLINIMIX/DEXTROSE (5/15) ...	130
cefoxitin sodium.....	130	CLINIMIX/DEXTROSE (5/20) ...	130
ceftriaxone sodium injection solution reconstituted 1 gm, 2 gm, 250 mg, 500 mg	130	CLINIMIX/DEXTROSE (5/25) ...	130
ceftriaxone sodium intravenous solution reconstituted 10 gm.	130	CLINISOL SF.....	130
cefuroxime sodium injection solution reconstituted 7.5 gm, 750 mg.....	130	colistimethate sodium (cba)	130
cefuroxime sodium intravenous solution reconstituted 1.5 gm	130	COMETRIQ (100 MG DAILY DOSE)	18
chlorpromazine hcl oral.....	130	COMETRIQ (140 MG DAILY DOSE)	18
chlorpropamide.....	51	COMETRIQ (60 MG DAILY DOSE)	18
chlorzoxazone oral tablet 500 mg	50	COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE	19
CINRYZE.....	130	COTELLIC.....	20
ciprofloxacin in d5w intravenous solution 200 mg/100ml	130	cromolyn sodium inhalation	130
clemastine fumarate oral tablet 2.68 mg.....	41	cyclobenzaprine hcl oral	50
CLIMARA PRO.....	47	cyclophosphamide oral capsule	131
clindamycin phosphate injection solution 600 mg/4ml.....	130	cyclosporine modified.....	131
CLINIMIX E/DEXTROSE (2.75/10)	130	cyclosporine oral capsule	131
CLINIMIX E/DEXTROSE (2.75/5)	130	cyproheptadine hcl oral	41
CLINIMIX E/DEXTROSE (4.25/10)	130	D	
CLINIMIX E/DEXTROSE (4.25/25)	130	dalfampridine er	7
CLINIMIX E/DEXTROSE (4.25/5)	130	daptomycin intravenous solution reconstituted 500 mg	131
CLINIMIX E/DEXTROSE (5/15) .	130	DEPO-PROVERA INTRAMUSCULAR SUSPENSION 400 MG/ML.....	131
CLINIMIX E/DEXTROSE (5/20) .	130	dextrose intravenous solution 10 % , 5 %	131
CLINIMIX E/DEXTROSE (5/25) .	130	dextrose-nacl intravenous solution 10-0.2 % , 10-0.45 % , 2.5-0.45 % , 5-0.2 % , 5-0.225 % , 5-0.33 % , 5-0.45 % , 5-0.9 %	131
CLINIMIX/DEXTROSE (2.75/5) .	130	diclofenac sodium transdermal gel 1 % , 3 %.....	21
CLINIMIX/DEXTROSE (4.25/10)	130	diclofenac sodium transdermal solution	21
CLINIMIX/DEXTROSE (4.25/20)	130	DIGITEK ORAL TABLET 250 MCG	39
CLINIMIX/DEXTROSE (4.25/25)	130	DIGOX ORAL TABLET 250 MCG..	39
		digoxin oral solution.....	39
		digoxin oral tablet 250 mcg.....	39
		diphtheria-tetanus toxoids dt ..	131
		dipyridamole oral.....	48
		disopyramide phosphate oral.....	39

HealthSun Health Plans
2019 Prior Authorization Criteria

doripenem intravenous solution
reconstituted 500 mg 131
doxercalciferol oral 131
DOXY 100 131
dronabinol..... 131
duramorph 131

E

ELIGARD..... 131
ENDARI22
ENGERIX-B INJECTION
SUSPENSION 10 MCG/0.5ML, 20
MCG/ML 131
entecavir34
ENVARBUS XR 131
ERAXIS..... 131
ERLEADA23
ERYTHROCIN LACTOBIONATE
INTRAVENOUS SOLUTION
RECONSTITUTED 500 MG 131
ESBRIET24
estradiol oral47
estropipate oral tablet 0.75 mg ..47

F

FARYDAK26
FENTORA BUCCAL TABLET 100
MCG27
FIRMAGON 131
FLEBOGAMMA DIF INTRAVENOUS
SOLUTION 5 GM/50ML 131
fluconazole in sodium chloride
intravenous solution 200-0.9
mg/100ml-%, 400-0.9
mg/200ml-% 131

FORTEO SUBCUTANEOUS
SOLUTION 600 MCG/2.4ML.....28
FREAMINE HBC 131
furosemide injection 131

G

GAMMAGARD INJECTION
SOLUTION 2.5 GM/25ML 131
GAMMAGARD S/D LESS IGA 131
GAMMAKED INJECTION SOLUTION
1 GM/10ML..... 131

GAMMAPLEX INTRAVENOUS
SOLUTION 10 GM/100ML, 10
GM/200ML, 20 GM/200ML, 5
GM/50ML..... 131

GAMUNEX-C INJECTION SOLUTION
1 GM/10ML 131
GENGRAF ORAL CAPSULE 100 MG,
25 MG 131
GENGRAF ORAL SOLUTION 131
gentamicin sulfate injection

solution 40 mg/ml..... 131
GILENYA ORAL CAPSULE 0.5 MG 29
GILOTRIF 30
glatiramer acetate 19
glyburide micronized 51
glyburide oral..... 51
glyburide-metformin 51
GOCOVRI 31
granisetron hcl oral..... 131
guanfacine hcl er 37

H

heparin sodium (porcine) injection
solution 1000 unit/ml, 10000
unit/ml, 20000 unit/ml, 5000
unit/ml..... 131
HEPATAMINE 131
HETLIOZ 36
HP ACTHAR..... 1

I

IBRANCE 53
ICLUSIG 54
IDHIFA..... 55
IMBRUVICA 56
imipenem-cilastatin intravenous
solution reconstituted 250 mg
..... 131
IMOVAX RABIES 131
indomethacin er 38
indomethacin oral 38
INLYTA..... 57
INTRAROSA 58
INTRON A..... 131
INVANZ INJECTION..... 59
IONOSOL-MB IN D5W 131

HealthSun Health Plans
2019 Prior Authorization Criteria

ipratropium bromide inhalation 131
ipratropium-albuterol..... 131
IRESSA.....60
ISOLYTE-P IN D5W..... 131
ISOLYTE-S 131

J

JAKAFI61
JUXTAPID ORAL CAPSULE 10 MG,
20 MG, 5 MG 62, 63

K

KALYDECO64
kcl in dextrose-nacl intravenous
solution 10-5-0.45 meq/l-%-%,
20-5-0.2 meq/l-%-%, 20-5-0.33
meq/l-%-%, 20-5-0.45 meq/l-
-%-%, 20-5-0.9 meq/l-%-%, 30-
5-0.45 meq/l-%-%, 40-5-0.45
meq/l-%-%, 40-5-0.9 meq/l-%-
% 131

kcl-lactated ringers-d5w..... 131
ketorolac tromethamine oral 38
KISQALI 200 DOSE65
KISQALI 400 DOSE65
KISQALI 600 DOSE65
KISQALI FEMARA 200 DOSE65
KISQALI FEMARA 400 DOSE65
KISQALI FEMARA 600 DOSE65
KORLYM.....66
KYNAMRO SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE
..... 67, 68

L

LAZANDA27
LENVIMA 10 MG DAILY DOSE.....69
LENVIMA 14 MG DAILY DOSE.....69
LENVIMA 18 MG DAILY DOSE.....69
LENVIMA 20 MG DAILY DOSE.....69
LENVIMA 24 MG DAILY DOSE.....69
LENVIMA 8 MG DAILY DOSE69
LETAIRIS70
LEUKINE INTRAVENOUS..... 131
leuprolide acetate injection 131
levalbuterol hcl inhalation
nebulization solution 0.31

mg/3ml, 0.63 mg/3ml, 1.25
mg/0.5ml, 1.25 mg/3ml 131
levocarnitine oral solution 131
levocarnitine oral tablet..... 131
levofloxacin in d5w intravenous
solution 500 mg/100ml, 750
mg/150ml..... 132
levofloxacin intravenous 132
lidocaine external patch 5 % 71
linezolid intravenous solution 600
mg/300ml..... 132
LONSURF 72
LUPRON DEPOT (1-MONTH) .73, 74
LUPRON DEPOT (3-MONTH) .73, 74
LUPRON DEPOT (4-MONTH) .73, 74
LUPRON DEPOT (6-MONTH) .73, 74
LYNPARZA 75

M

magnesium sulfate injection
solution 50 %, 50 % (10ml
syringe)..... 132
MATULANE 76
MAVYRET 35
megestrol acetate oral suspension
625 mg/5ml..... 46
MEKINIST..... 77
MENEST ORAL TABLET 0.3 MG,
0.625 MG, 1.25 MG 47
meperidine hcl injection solution
100 mg/ml, 25 mg/ml, 50 mg/ml
..... 38
meperidine hcl oral 38
meropenem 132
methocarbamol oral 50
methotrexate oral..... 132
methotrexate sodium (pf) injection
solution 50 mg/2ml..... 132
methotrexate sodium injection
solution 250 mg/10ml 132
methyldopa oral 42
methyldopa-hydrochlorothiazide 42
methylphenidate hcl er (cd) oral
capsule extended release 10 mg,
20 mg, 40 mg, 50 mg, 60 mg. 37

HealthSun Health Plans
2019 Prior Authorization Criteria

methylphenidate hcl er oral tablet
extended release 20 mg37
methylphenidate hcl er oral tablet
extended release 24 hour37
methylphenidate hcl oral37
metronidazole in nacl intravenous
solution 500-0.79 mg/100ml-%
..... 132
MIRCERA INJECTION SOLUTION
PREFILLED SYRINGE 100
MCG/0.3ML 132
modafinil91
moxifloxacin hcl in nacl 132
MYCAMINE 132
mycophenolate mofetil..... 132
mycophenolate sodium 132

N

nafcillin sodium injection solution
reconstituted 1 gm 132
nafcillin sodium intravenous
solution reconstituted 10 gm. 132
NEBUPENT 132
NEPHRAMINE..... 132
NERLYNX 79
NEXAVAR80
nifedipine oral.....45
NINLARO81
nitrofurantoin macrocrystal oral .52
nitrofurantoin monohyd macro...52
NORDITROPIN FLEXPRO..... 32, 33
NORMOSOL-M IN D5W..... 132
NORMOSOL-R IN D5W 132
NORMOSOL-R PH 7.4..... 132
NORTHERA.....82
NUEDEXTA83
NUPLAZID ORAL TABLET 17 MG .84
nutrilipid intravenous emulsion 20
% 132

O

OCTAGAM INTRAVENOUS
SOLUTION 1 GM/20ML, 2
GM/20ML 132
octreotide acetate injection
solution 100 mcg/ml, 1000

mcg/ml, 200 mcg/ml, 50
mcg/ml, 500 mcg/ml..... 132
ondansetron 132
ondansetron hcl oral 132
OPSUMIT 85
ORKAMBI ORAL TABLET..... 86
orphenadrine citrate er..... 50
OSPHENA 87

P

paricalcitol oral..... 132
penicillin g potassium injection
solution reconstituted 20000000
unit 132
penicillin g sodium 132
PENTAM 132
pentazocine-naloxone hcl..... 38
perphenazine oral tablet 4 mg, 8
mg 132
piperacillin sod-tazobactam so
intravenous solution
reconstituted 2.25 (2-0.25) gm,
3.375 (3-0.375) gm, 4.5 (4-0.5)
gm 132
PLASMA-LYTE 148 132
PLASMA-LYTE A..... 132
PLEGRIDY..... 78
PLEGRIDY STARTER PACK
SUBCUTANEOUS SOLUTION PEN-
INJECTOR 78
PLENAMINE 132
POMALYST..... 90
potassium chloride in dextrose
intravenous solution 20-5 meq/l-
%, 40-5 meq/l-% 132
potassium chloride in nacl
intravenous solution 20-0.45
meq/l-%, 20-0.9 meq/l-% ... 132
potassium chloride intravenous
solution 2 meq/ml, 20
meq/100ml 132
PRALUENT SUBCUTANEOUS
SOLUTION PEN-INJECTOR.88, 89
PREMARIN ORAL..... 47
PREMASOL..... 132

HealthSun Health Plans
2019 Prior Authorization Criteria

PREMPRO.....47
 PRIVIGEN INTRAVENOUS
 SOLUTION 20 GM/200ML..... 132
 PROCALAMINE..... 132
 prochlorperazine maleate oral .. 132
 PROCRIT.....25
 PROLASTIN-C INTRAVENOUS
 SOLUTION RECONSTITUTED
 1000 MG 132
 promethazine hcl oral syrup.....40
 promethazine hcl oral tablet40
 promethazine hcl rectal.....40
 promethazine vc plain oral solution
 41
 PROSOL..... 132
 PULMOZYME..... 132
R
 RABAVERT 132
 RAPAMUNE ORAL SOLUTION.... 132
 RECOMBIVAX HB..... 132
 REGRANEX.....92
 REPATHA 88, 89
 REPATHA PUSHTRONEX SYSTEM
 88, 89
 REPATHA SURECLICK 88, 89
 rifampin intravenous..... 132
 RUBRACA..... 93, 94
 RYDAPT95
S
 SAMSCA96
 SENSIPAR 132
 sildenafil citrate oral tablet 20 mg
 97
 sirolimus oral..... 132
 sodium chloride injection solution
 2.5 meq/ml 132
 sodium chloride intravenous
 solution 0.45 %, 0.9 %, 3 %, 5
 % 133
 sodium lactate intravenous solution
 5 meq/ml 133
 SPRYCEL.....98
 STIVARGA.....99
 SUTENT 100

SYLATRON SUBCUTANEOUS KIT
 200 MCG, 300 MCG, 600 MCG
 101
 SYMDEKO..... 102
 SYNDROS..... 133
 SYNRIPO..... 103
T
 tacrolimus oral 133
 TAFINLAR..... 105
 TAGRISSO..... 106
 TARCEVA..... 107
 TASIGNA 108
 TECFIDERA 109
 TEFLARO 133
 TENIVAC 133
 tetanus-diphtheria toxoids td .. 133
 tigecycline 133
 TOBI PODHALER..... 110
 tobramycin inhalation..... 133
 tobramycin sulfate injection
 solution 10 mg/ml, 80 mg/2ml
 133
 TPN ELECTROLYTES INTRAVENOUS
 SOLUTION 133
 TRACLEER 111
 TRAVASOL..... 133
 TRELSTAR MIXJECT 133
 tretinoin external cream 112
 tretinoin external gel 0.01 %,
 0.025 % 112
 TREXALL 133
 trientine hcl 104
 trihexyphenidyl hcl 43
 TROPHAMINE INTRAVENOUS
 SOLUTION 10 %..... 133
 TWINRIX 133
 TYKERB..... 113
 TYMLOS 114
U
 UPTRAVI 115
V
 vancomycin hcl intravenous
 solution reconstituted 10 gm,
 1000 mg, 500 mg 133

HealthSun Health Plans
2019 Prior Authorization Criteria

VARUBI ORAL 133
 VENCLEXTA 116
 VENCLEXTA STARTING PACK ... 116
 VERZENIO 117
 VOSEVI 35
 VOTRIENT 118
X
 XALKORI 119
 XATMEP 133
 XTANDI 120
 XURIDEN 121
Y
 YONSA 122

Z
 zaleplon 49
 ZEJULA 123
 ZELBORAF 124
 ZOLINZA 125
 zolpidem tartrate er 49
 zolpidem tartrate oral 49
 ZORTRESS ORAL TABLET 0.25 MG,
 0.5 MG, 0.75 MG 133
 ZYDELIG 126, 127
 ZYKADIA 128
 ZYTIGA 129